EDITORIAL

Laboratory medicine: Closer clinical collaboration will lead to evidence-based reporting of results

Amro Maarouf, Nathan Lorde, Mark Livingston, Sudarshan Ramachandran

When using biochemical analytes in the management of patients, the value of the analyte is often considered in relation to the reference range calculated from the distribution of the analyte in a healthy cohort [1]. Such distributions are either symmetrically (Gaussian) or asymmetrically distributed. The middle 95% values of the healthy cohort distribution are utilized as the reference range assumes that the reference cohort is an apt comparison group for the specific pathology. For example, the measurement of cortisol provides an important reminder of the pitfalls of reference range dependence. Although most clinicians are aware of this hormone's diurnal pattern, the empirical undertaking of an early morning basal sample in order to assess the integrity of the hypothalamic-pituitary-adrenal axis may in certain circumstances be inappropriate, such as in individuals with altered sleep-wake cycle. Furthermore, a numeral value that straddles the higher end of a

```
Amro Maarouf<sup>1</sup>, Nathan Lorde<sup>1</sup>, Mark Livingston<sup>2,3</sup>, Sudarshan Ramachandran<sup>1,4,5</sup>
```

<u>Affiliations:</u> ¹Department of Clinical Biochemistry, University Hospitals Birmingham NHS Foundation Trust, Rectory Road, Sutton Coldfield, West Midlands, B75 7RR, United Kingdom; ²Department of Clinical Biochemistry, Black Country Pathology Services, Walsall Manor Hospital, Walsall, United Kingdom; ³School of Medicine and Clinical Practice, Faculty of Science & Engineering, The University of Wolverhampton, United Kingdom; ⁴University Hospitals of North Midlands/Faculty of Health Sciences, Staffordshire University/Institute of Science and Technology, Keele University, Staffordshire, United Kingdom; ⁵College of Engineering, Design & Physical Sciences, Brunel University London, Uxbridge, United Kingdom.

<u>Corresponding Author:</u> Professor Sudarshan Ramachandran, Department of Clinical Biochemistry, University Hospitals Birmingham NHS Foundation Trust, Good Hope Hospital, Rectory Road, Sutton Coldfield, West Midlands B75 7RR, United Kingdom; Email: sud.ramachandran@ heartofengland.nhs.uk

Received: 23 April 2021 Published: 20 May 2021 reported reference range—in an inpatient setting—may falsely reassure a clinician into believing there is adequate adrenal reserve, when actually representing an adrenal crisis. It is therefore imperative that an assayed sample is taken in the context of a thorough understanding of the wider medical and psychosocial setting.

Thus, it is important to evaluate as to what the reference range adds to the clinical decision-making process and whether it should be replaced by action limits. While knowledge of this distribution can be invaluable, it must be used in conjunction with the clinical phenotype while diagnosing a condition, as well as the available evidence base when making treatment decisions. We ask the question as to whether the value obtained from a biochemistry test is perhaps better when compared with both the distributions of healthy individuals and those with the specific pathology. It is further complicated by heterogeneity existing within most disease states, which may require knowledge of differing distributions. Many clinicians can intuitively bypass these issues through innate awareness and experience. They are able to judge whether an abnormal biochemical test is clinically relevant or not, regardless of whether it lies within or outside the normal distribution. Importantly, many medical schools now devote little education in describing the derivation of reference ranges and the impact of heterogeneity. Hence, biochemical tests and reference ranges are often considered by trainee doctors as absolute predictors of health or disease, leading to a deficient practice of medicine. But we know that, even in a "normal" population, statistically speaking, a test result will lie outside the reference range in 5% of cases (1 in 20)! This is perhaps why the term "reference range" is preferential to "normal range." We now provide two clinical examples; one where reference ranges have been successfully superseded by evidence-based action limits (dyslipidemia), and another where confusion exists with very different reference ranges quoted and no action limits reported (male adult-onset hypogonadism).

Most clinical biochemistry laboratories do not report lipid distributions in healthy subjects. For example the Mayo Clinic (https://www.mayoclinic.org/diseasesconditions/high-blood-cholesterol/diagnosis-treatment/ drc-20350806; accessed on 16th April 2021) categorizes

Edorium J Biochem 2021;5:100007B01AM2021. *www.edoriumjournalofbiochemistry.com*

total cholesterol, low density lipoprotein (LDL)cholesterol, high density lipoprotein cholesterol, and triglyceride concentrations in relation to heart disease in a patient friendly guideline format. LDL-cholesterol levels are issued, as follows, in relation to the likelihood of heart disease risk.

- <1.8 mmol/L (70 mg/dL): best for people who have heart disease or diabetes.
- <2.6 mmol/L (<100 mg/dL): optimal for people at risk of heart disease.
- 2.6–3.3 mmol/L (100–129 mg/dL): near optimal if there is no heart disease, high if there is heart disease.
- 3.4–4.1 mmol/L (130–159 mg/dL): borderline high if there is no heart disease, high if there is heart disease.
- 4.1–4.9 mmol/L (160–189 md/dL): high if there is no heart disease, very high if there is heart disease.
- >4.9 mmol/L (190 mg/dL): very high.

However, the above guidance does not completely mirror the recommendations of the American Heart Association/American College of Cardiology (AHA/ACC) issued in 2018, especially regarding the use of risk scores in primary prevention [2]. Interestingly, the 2019 European Society of Cardiology (ESC)/European Atherosclerosis Society (EAS) guidelines, issued in 2019, significantly differ from those issued by the AHA/ ACC [3]. In the United Kingdom (UK), the information of lipid levels by the National Health Service (https:// www.nhs.uk/conditions/high-cholesterol/cholesterollevels/; accessed on 16th April 2021) is lagging behind current trial evidence and the ESC/EAS guidelines. Thus, it is not sufficient in a field such as lipidology with constantly advancing evidence, to just adopt action limits, but rather to periodically update them through the continued review of the latest clinical trials/guidelines by international societies. Furthermore, heterogeneity must also be considered, as the lipid thresholds will differ between various dyslipidemia subgroups (e.g., familial hypercholesterolemia and metabolic syndrome) and therefore should also be highlighted in any reports [4].

Adult-onset hypogonadism in males is defined by low testosterone concentrations and associated symptoms [5]. Identifying this clinical phenotype is important as it has a high prevalence (6–12% in males and about 40% in type 2 diabetes) as well as being associated with increased morbidity and mortality [5, 6]. A recent audit by our group in the UK showed male testosterone reference ranges varying significantly between laboratories [7]. Furthermore, no useful clinical guidance was provided in most of the reports, with only a few quoting published action limits. There is mounting evidence regarding the benefits of testosterone therapy in men with adult-onset hypogonadism [5], and it is essential that these should be communicated by the laboratories, to the service users.

The above examples demonstrate how laboratories report analytes in two differing pathologies; one with mature evidence (dyslipidemia) and the other with evidence in its infancy (adult-onset hypogonadism). Even in the former, it is clear that reports issued by the laboratory must include the ever-increasing evidence. In the latter, the laboratory has an obligation, not just to harmonize reference ranges, but to monitor evidence and move the reporting from reference ranges toward action limits, and also educate the requesting clinicians from a myriad of specialties (urology, endocrinology, metabolic physicians, and general practice) in the appropriate management of these patients. The key to the optimization of reporting/education is close collaboration with specialists in all disciplines in which management is influenced by laboratory measurements of analytes.

Keywords: Adult-onset hypogonadism, Dyslipidemia guidelines, Reference ranges

How to cite this article

Maarouf A, Lorde N, Livingston M, Ramachandran S. Laboratory medicine: Closer clinical collaboration will lead to evidence-based reporting of results. Edorium J Biochem 2019;4:100007B01AM2021.

Article ID: 100007B01AM2021

doi: 10.5348/100007B01AM2021ED

REFERENCES

- Ramachandran S, Konig CS, Hackett G, Livingston M, Strange RC. Managing clinical heterogeneity: An argument for benefit based action limits. Journal of Medical Diagnostics and Therapy 2018;1:034701.
- 2. Grundy SM, Stone NJ. 2018 American Heart Association/American College of Cardiology Multisociety Guideline on the Management of Blood Cholesterol: Primary Prevention. JAMA Cardiol 2019;4(5):488–9.
- 3. Singh M, McEvoy JW, Khan SU, et al. Comparison of transatlantic approaches to lipid management: The AHA/ACC/Multisociety Guidelines vs the ESC/EAS Guidelines. Mayo Clin Proc 2020;95(5):998–1014.
- 4. Shipman KE, Strange RC, Ramachandran S. Use of fibrates in the metabolic syndrome: A review. World J Diabetes 2016;7(5):74–88.
- 5. Hackett G, Kirby M, Edwards D, et al. British Society for Sexual Medicine guidelines on adult testosterone deficiency, with statements for UK practice. J Sex Med 2017;14(12):1504–23.

Edorium J Biochem 2021;5:100007B01AM2021. *www.edoriumjournalofbiochemistry.com*

- 6. Pye SR, Huhtaniemi IT, Finn JD, et al. Late-onset hypogonadism and mortality in aging men. J Clin Endocrinol Metab 2014;99(4):1357–66.
- Livingston M, Downie P, Hackett G, Marrington R, Heald A, Ramachandran S. An audit of the measurement and reporting of male testosterone levels in UK clinical biochemistry laboratories. Int J Clin Pract 2020;74(111):e1360.

Author Contributions

Amro Maarouf – Conception of the work, Design of the work, Drafting the work, Revising the work critically for important intellectual content, Final approval of the version to be published, Agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

Nathan Lorde – Conception of the work, Design of the work, Drafting the work, Revising the work critically for important intellectual content, Final approval of the version to be published, Agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

Mark Livingston – Conception of the work, Design of the work, Revising the work critically for important intellectual content, Final approval of the version to be published, Agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved Sudarshan Ramachandran – Conception of the work, Design of the work, Drafting the work, Final approval of the version to be published, Agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

Guarantor of Submission

The corresponding author is the guarantor of submission.

Source of Support

None.

Consent Statement

Written informed consent was obtained from the patient for publication of this article.

Conflict of Interest

Authors declare no conflict of interest.

Data Availability

All relevant data are within the paper and its Supporting Information files.

Copyright

© 2021 Amro Maarouf et al. This article is distributed under the terms of Creative Commons Attribution License which permits unrestricted use, distribution and reproduction in any medium provided the original author(s) and original publisher are properly credited. Please see the copyright policy on the journal website for more information.











Submit your manuscripts at

www.edoriumjournals.com







EDORIUM JOURNAL OF





JOURNAL OF CASE REPORTS AND IMAGES IN PATHOLOGY







PSYCHIATRY

EDORIUM JOURNAL OF

ANESTHESIA

JOURNAL OF CASE REPORTS AND IMAGES IN ORTHOPEDICS AND RHEUMATOLOGY







EDORIUM JOURNAL OF

INTERNATIONAL JOURNAL OF CASE REPORTS AND IMAGES





VIDEO JOURNAL OF BIOMEDICAL SCIENCE





INTERNATIONAL JOURNAL OF BLOOD TRANSFUSION AND IMMUNOHEMATOLOGY



